

CertiFeed HV

22-20 SF OTC CALF MILK REPLACER MEDICATED

Each pound of this milk replacer contains 10 mg. of oxytetracycline. Feed continuously to calves up to 250 lbs. Feed at a rate to provide each calf 0.05 to 0.1 mg of oxytetracycline per lb. of bodyweight daily.

Milk Replacer to be fed to herd replacement calves.

For calves (up to 250 lbs.) for increased rate of gain and improved feed efficiency.

ACTIVE DRUG INGREDIENT(S)	
Oxytetracycline	20 g/ton
GUARANTEED ANALYSIS	
Crude Protein , Min	22.00 %
Crude Fat , Min	20.00 %
Crude Fiber , Max	0.75 %
Calcium , Min	0.75 %
Calcium , Max	1.25 %
Phosphorus , Min	0.70 %
Vitamin A , Min	30000 IU/lb.
Vitamin D3 , Min	10000 IU/lb.
Vitamin E , Min	150 IU/lb.
Ascorbic Acid , Min	100.00 mg./lb.
Vitamin B-12 , Min	0.080 mg./lb.

INGREDIENTS
Dried whey, soy flour, animal fat (preserved with BHA, BHT, citric acid & ethoxyquin), dried whey product, dried whey protein concentrate, calcium carbonate, dried skimmed milk, lecithin, L-lysine, sodium silico aluminate, maltodextrin, DL-methionine, ethoxylated mono-diglycerides, propylene glycol, ferrous sulfate, vitamin E supplement, magnesium sulfate, choline chloride, artificial flavor, ascorbic acid, selenium yeast, brewer's dried yeast, zinc sulfate, vitamin A supplement, niacin supplement, manganese sulfate, vitamin D3 supplement, copper sulfate, calcium pantothenate, biotin, riboflavin supplement, menadione sodium bisulfite complex (source of vitamin K activity), vitamin B12 supplement, thiamine mononitrate, pyridoxine hydrochloride, silicon dioxide, calcium iodate, folic acid, cobalt sulfate.

Important: Follow feeding directions.

Manufactured for:
Cooperative Feed Dealers
Conklin, NY 13748

NET WEIGHT 50 LB (22.68 kg)
01-9150

MIXING DIRECTIONS
Enclosed in each bag is a cup that holds 10 ounces of dry milk replacer powder level full. Mix milk replacer powder with warm (110-115°F) water. For accurate feeding, measure cup weight periodically.

Feed high quality colostrum for the first day, even if your calf is more than 24 hours old. Begin feeding milk replacer on day 2.

Add 1 enclosed cup, level full, of milk replacer powder to 2 quarts of warm water (110-115°F) and mix thoroughly. For large batches, mix 10 full cups (6 1/4 lbs.) of milk replacer powder to 5 gallons of warm water.

FEEDING DIRECTIONS
Large breeds: Feed 2 quarts reconstituted milk replacer twice daily.

Small breeds: Feed 1 1/2 to 2 quarts reconstituted milk replacer twice daily.

OFFER FRESH WATER AT ALL TIMES

- CALF CARE RECOMMENDATIONS**
- 1) Feed large breed calves 4 quarts of high quality colostrum immediately after birth, within 2 hours if possible, and again 12 hours later. Adjust amounts for small calves. If little or no maternal colostrum is available, feed a supplement.
 - 2) Begin feeding milk replacer on day 2.
 - 3) Begin feeding calf starter free-choice during the first week. Feed a small amount at first and replace with fresh feed daily.
 - 4) Provide fresh, clean water at all times.
 - 5) Wean the healthy calf from milk replacer when its intake of calf starter exceeds 2 pounds daily during at least 3 consecutive days.
 - 6) Stressful environmental conditions such as cold temperatures, may necessitate a change in the feeding rate of milk replacer. Contact your nutritionist for specific instructions.

While this product is nutritionally complete, best results are achieved when calves are fed an all milk protein calf milk replacer from the second day of life, then transitioned gradually to this product after 3 weeks of age.