

CertiFeed

20-20 AM BOV CALF MILK REPLACER MEDICATED

Milk Replacer to be fed to herd replacement calves. Control of coccidiosis caused by Eimeria bovis and Eimeria zuernii in cattle.

ACTIVE DRUG INGREDIENT(S)
Lasalocid ... 72 g/ton

GUARANTEED ANALYSIS	
Crude Protein	, Min 20.00 %
Crude Fat	, Min 20.00 %
Crude Fiber	, Max 0.15 %
Calcium	, Min 0.75 %
Calcium	, Max 1.25 %
Phosphorus	, Min 0.70 %
Vitamin A	, Min 30000 IU/lb.
Vitamin D3	, Min 10000 IU/lb.
Vitamin E	, Min 150 IU/lb.
Ascorbic Acid	, Min 100.00 mg./lb.
Vitamin B-12	, Min 0.080 mg./lb.

INGREDIENTS
Dried whey product, dried whey protein concentrate, dried whey, animal fat (preserved with BHA, BHT, citric acid & ethoxyquin), calcium carbonate, dried skimmed milk, sodium silico aluminate, maltodextrin, ferrous sulfate, vitamin E supplement, L-lysine, magnesium sulfate, choline chloride, artificial flavor, DL-methionine, ascorbic acid, selenium yeast, brewer's dried yeast, zinc sulfate, vitamin A supplement, lecithin, ethoxylated mono-diglycerides, propylene glycol, niacin supplement, manganese sulfate, vitamin D3 supplement, copper sulfate, calcium pantothenate, biotin, menadione sodium bisulfite complex (source of vitamin K activity), riboflavin supplement, vitamin B12 supplement, thiamine mononitrate, pyridoxine hydrochloride, calcium iodate, folic acid, silicon dioxide, cobalt sulfate.

Important: Follow feeding directions.

*** * C A U T I O N : * ***
The safety of Lasalocid in unapproved species has not been established. Do not allow horses or other equines, dogs or cats, access to Lasalocid as ingestion may be fatal.

*** * W A R N I N G : * ***
A withdrawal period has not been established for Lasalocid in pre-ruminating cattle. Do not use in calves to be processed for veal.

Feed to provide each calf 1 mg. of lasalocid per 2.2 lb.(1 kg.) of body weight per day. Feed to cattle weighing up to 800 lb. with a maximum of 360 mg/head/day. One pound of milk replacer powder will medicate 79.2 pounds of bodyweight.

MIXING DIRECTIONS
Enclosed in each bag is a cup that holds 10 ounces of dry milk replacer powder level full. Mix milk replacer powder with warm (110-115°F) water. For accurate feeding, measure cup weight periodically.

Feed high quality colostrum for the first day, even if your calf is more than 24 hours old. Begin feeding milk replacer on day 2.

Add 1 enclosed cup, level full, of milk replacer powder to 2 quarts of warm water (110-115°F) and mix thoroughly. For large batches, mix 10 full cups (6 1/4 lbs.) of milk replacer powder to 5 gallons of warm water.

FEEDING DIRECTIONS
Large breeds: Feed 2 quarts reconstituted milk replacer twice daily.

Small breeds: Feed 1 1/2 to 2 quarts reconstituted milk replacer twice daily.

IMPORTANT: Mix milk replacer powder into water and feed immediately. When preparing large batches keep the mixture agitated while dispensing into individual feeding containers to prevent settling, and thus the feeding of incorrect lasalocid levels.

OFFER FRESH WATER AT ALL TIMES

- CALF CARE RECOMMENDATIONS**
- 1) Feed large breed calves 4 quarts of high quality colostrum immediately after birth, within 2 hours if possible, and again 12 hours later. Adjust amounts for small calves. If little or no maternal colostrum is available, feed a supplement.
 - 2) Begin feeding milk replacer on day 2.
 - 3) Begin feeding calf starter free-choice during the first week. Feed a small amount at first and replace with fresh feed daily.
 - 4) Provide fresh, clean water at all times.
 - 5) Wean the healthy calf from milk replacer when its intake of calf starter exceeds 2 pounds daily during at least 3 consecutive days.
 - 6) Stressful environmental conditions such as cold temperatures, may necessitate a change in the feeding rate of milk replacer. Contact your nutritionist for specific instructions.

Manufactured for:
Cooperative Feed Dealers
Conklin, NY 13748

NET WEIGHT 50 LB (22.68 kg)

01-9143

REV. 0309